

Patients with TBI or blow to the head With LOC on Warfarin or Therapeutic Heparin/LMWH

Patients with TBI or blow to the head With LOC on oral anticoagulants

Initial CT negative and:
INR > 1.5
or
PTT > 50
or
Factor Xa < 40%

Initial CT positive and
INR > 1.5
or
PTT > 50
or
Factor Xa < 40 %

Initial CT positive

Initial CT Negative and
TT > 30 sec
INR > 1.5

- Hold medication
- Observe Pt. for at least 6 hours in hospital
- Repeat CT in 12-24 hrs (6 hrs. if significant injury)
- If INR > 10 hold warfarin and give vitamin K (see procedure PROC_CM_A-30F)
- If reversed:
- Observe Pt. for at least 24 hours in hospital

- Hold Medication
- Admit Pt. for at least 24 hours in hospital
- Acutely reverse anticoagulation (See procedure PROC_CM_A-30(a-f))

- Hold Medication
- Admit Pt. in hospital for at least 24 hours
- Obtain repeat Head CT (6 hrs. and 24 Hrs. if no change on first repeat)
- Acutely Reverse (See PROC_CM_A-30C)
- Hematology consult
- If Pt. is stable consider:
 - Hemodialysis
 - Plasmapheresis

- Hold Medication
- Observe patient in hospital for at least 24 hrs
- Repeat CT in 12-24 hours (6 hours if significant injury)