

Information for You from Your Health Care Team

Nulytely Colonoscopy Preparation (3 Day Prep)

Your colonoscopy will be performed by: _____

Date and Time of Colonoscopy: _____ Arrive at: _____.

Your procedure will be done in the Endoscopy Suite at:

Hill Medical Building Suite 206

University Hospital, 750 E. Adams St.

The colon must be completely cleansed and empty. This is achieved by drinking the bowel cleansing solution given to you as indicated below. Failure to follow these instructions could lead to an unsuccessful colonoscopy that may have to be repeated.

Three days prior to the procedure:

- Begin a clear liquid diet and remain on a clear liquid diet until after your colonoscopy is completed.
- **Clear Liquid Diet**
 - **Soups:** Clear bouillon, chicken broth or vegetable broth.
 - **Beverages:** Tea and Coffee (without cream/milk), Kool-Aid and Gatorade (**no red or purple**), apple juice, cranberry juice, grape juice, and carbonated beverages.
 - **Dessert:** Italian ice, Popsicles, and Jell-O (**no red or purple**).
 - Drink 1 gallon of Nulytely on the 2nd day of clear liquids and 1gallon of Nulytely on the 3rd day of clear liquids. Start drinking the preparation between 2:30 PM and 4:30 PM. Drink 1 glass (8oz) of the Nulytely every 10-15 minutes until the gallon is finished. If you become nauseated, stop drinking the solution until the nausea subsides. Then restart the solution, leaving 20 minute intervals or longer between glassfuls. **It is very important that you finish the preparation without vomiting.**
 - This will cause you to have loose, watery bowel movements.
 - You may drink as much water as you like until midnight.
 - Nothing after midnight, except medications.
 - **If you need to cancel or change your appointment, please call the Endoscopy scheduler at 464-1600 48 hours in advance.**
- **Because of the medications given to you during the examination, You must have someone accompany you home after you are discharged. If you come without someone, your test will not be done.**
- **Medications:** Continue your usual medications (except for those listed below) as scheduled. The morning of your colonoscopy, you may take your essential medications such as those for blood pressure, heart, thyroid, and seizure with a small amount of water.
 - Notify your doctor **NOW** if you are taking any “blood thinning” medications such as **Coumadin (Warfarin) or Plavix.**
 - **For diabetics:** If you are a diabetic taking insulin injections, your dose may need to be adjusted for the three days you are on clear liquids. Please contact the physician who monitors your diabetes for instructions. Do not take your insulin the morning of your procedure. Unless told differently by your physician, continue your usual dose the day after the procedure.

- **Stop taking iron pills** 7 days prior to Colonoscopy.
- Please leave any valuables at home.

If you have any questions or need to cancel or change your appointment, please call University Gastroenterology at (315) 464-1600 or University Gastroenterology Endoscopy Suite at (315) 464-1616.